

## K-theory of operator algebras - written examination

- 5 1. Compute  $K_*(C(S^1))$  and  $K_*(S^2)$ .
- 5 2. Using the homotopy invariance of  $K$ -groups and remembering  $K_*(S^1)$  and  $K_*(S^2)$ , prove that  $S^2 \times S^1$  is not homotopic to  $S^3$ .
- 3 3. Using the functoriality of  $K$ -theory, prove that there is no retraction of the two-dimensional disc  $D$  to  $S^1$ .
- 4 4. For any action of  $\mathbb{Z}$  on the two-disc  $D$ , compute  $K_*(C(D) \rtimes \mathbb{Z})$ .
- 4 5. Let  $\mathcal{T}$  denote the Toeplitz algebra, and let  $s$  denote its generator (one-sided shift). Then there is a  $*$ -homomorphism  $\sigma : \mathcal{T} \rightarrow C(S^1)$ , given by  $\sigma(s) = u$ ,  $u$  the unitary generator of  $C(S^1)$ . Let  $\mathcal{T} \oplus_\sigma \mathcal{T} := \{(f, g) \in \mathcal{T} \oplus \mathcal{T} \mid \sigma(f) = \sigma(g)\}$  (pull-back). Taking for granted that  $K_0(\mathcal{T}) \simeq \mathbb{Z}$  is generated by  $[1]_0$ , but assuming no knowledge of  $K_1(\mathcal{T})$ , show that  $K_0(\mathcal{T} \oplus_\sigma \mathcal{T})$  contains  $\mathbb{Z}$  as a direct summand.
- 4 6. Give an example of a  $C^*$ -algebra  $A$  for which  $K_0(A) = 0$ ,  $K_1(A) \neq 0$ . Show that if  $K_0(A) = 0$ ,  $K_1(A) \neq 0$ , then  $A$  cannot be the  $C^*$ -algebra of a directed graph with finitely many vertices.