

Dirac Operators and Spectral Geometry

Written examination

Each problem is worth 5 points, out of a possible total of 30.

1. Let $\text{Cl}_{p,q} := \text{Cl}(\mathbb{R}^{p+q}, g)$ be the real Clifford algebra for the symmetric bilinear form $g(x, x) = x_1^2 + \cdots + x_p^2 - x_{p+1}^2 - \cdots - x_{p+q}^2$, and let Cl_{pq}^0 be its even subalgebra. Prove that $\text{Cl}_{p+1,q}^0 \simeq \text{Cl}_{q,p}$ as \mathbb{R} -algebras.

2. Let \mathcal{A} be the unital $*$ -algebra generated by elements x, y, z , and consider $p = \frac{1}{2} \begin{pmatrix} 1+z & x+iy \\ x-iy & 1-z \end{pmatrix} \in M_2(\mathcal{A})$. Prove that the relations $p = p^* = p^2$ imply that \mathcal{A} is a polynomial algebra of functions on \mathbb{S}^2 , by showing that

$$\begin{aligned} x^* &= x, & y^* &= y, & z^* &= z, \\ [x, y] &= [x, z] = [y, z] &= 0, \\ x^2 + y^2 + z^2 &= 1. \end{aligned}$$

3. Show that the distance function on \mathbb{S}^1 determined by the Dirac operator $\mathcal{D} = -i d/d\theta$ is the arc length of the circle.

4. The Dirac operator on the sphere \mathbb{S}^2 with its usual round metric has eigenvalues $\pm k$, with respective multiplicities $2k$, for $k = 1, 2, 3, \dots$. Let $\sigma_N(T)$ denote the sum of the N largest singular values of a compact operator T . If $s > 0$, prove that

$$L = \lim_{N \rightarrow \infty} \frac{\sigma_N(|\mathcal{D}|^{-s})}{\log N}$$

exists, with $0 < L < +\infty$, if and only if $s = 2$. Also, find this value of L .

5. A function or distribution u on \mathbb{R}^n is homogeneous of degree λ if rescaling by a factor of t changes u to $t^\lambda u$. Show that the Dirac delta distribution on \mathbb{R}^n , defined by $\langle \delta, f \rangle := f(0)$, is homogeneous of degree $-n$.

6. If M is a compact manifold, its space of smooth functions $C^\infty(M)$, with the topology of uniform convergence of all derivatives, is known to be a unital Fréchet $*$ -algebra. Prove that $C^\infty(M)$ is a pre- C^* -algebra, i.e., there is a C^* -algebra in which $C^\infty(M)$ is closed under holomorphic functional calculus.

Easy oral questions

E. Describe the real Clifford algebra $\text{Cl}(\mathbb{R}^2)$, where \mathbb{R}^2 has its standard Euclidean structure.

E. When are two unital algebras A and B Morita equivalent? If M is an even-dimensional compact manifold, explain why the algebras $C(M)$ and $\Gamma(M, \text{Cl}(M))$ are Morita equivalent.

E. Identify the spectrum of the Dirac operator $\mathcal{D} = -i d/d\theta$ on the circle \mathbb{S}^1 with its trivial spinor module $\mathcal{S} = C^\infty(\mathbb{S}^1)$.

E. What is a (unital) spectral triple? What are its data, and what conditions must be satisfied by these data?

E. Given a positive elliptic pseudodifferential operator A of order $(-n)$ on a compact manifold of dimension n , why is $\text{Tr}_\omega A$ independent of the choice of Dixmier trace Tr_ω , and how does one compute $\text{Tr}_\omega A$?

E. Give an example of an isospectral deformation of a spectral triple.

More difficult oral questions

D. Let V be a Euclidean space of real dimension n . Explain why the *centre* $Z(\text{Cl}(V))$ of the Clifford algebra $\text{Cl}(V)$ has dimension 1 if n is even, and has dimension 2 if n is odd.

D. Describe briefly the construction of the irreducible representation of the complex Clifford algebra $\text{Cl}(\mathbb{R}^{2m})$ on the vector space underlying a complex exterior algebra.

D. Explain why the spinor bundle over the 3-sphere, $S \rightarrow \mathbb{S}^3$, is a trivial vector bundle.

D. Explain why the spinor bundle over the 2-sphere, $S \rightarrow \mathbb{S}^2$, is a trivial vector bundle.

D. Explain how one can obtain the area of the sphere \mathbb{S}^2 , knowing the spectrum of the Dirac operator for the round metric on \mathbb{S}^2 .

D. Suppose that (M, g) is a compact Riemannian spin manifold whose scalar curvature s is positive at each point of M . Outline how the Lichnerowicz formula $\mathcal{D}^2 = \Delta^S + \frac{1}{4}s$ allows one to prove that the Dirac operator has kernel $\{0\}$.

D. Outline the construction of the Dixmier traces as functionals on positive compact operators.

D. A compact operator T on a Hilbert space \mathcal{H} is called *measurable* if it belongs to the Dixmier trace class $\mathcal{L}^{1+}(\mathcal{H})$, and if the value of any $\text{Tr}_\omega T$ is independent of which Dixmier trace Tr_ω is used. How can one guarantee that an operator $T \in \mathcal{L}^{1+}(\mathcal{H})$ is measurable? If it is measurable, describe how to compute $\text{Tr}_\omega T$.

D. Describe in detail the definition of the Wodzicki residue, $\text{Wres } P$, of a classical pseudodifferential operator P on a compact manifold M .

D. If \mathcal{D} is the Dirac operator on an n -dimensional compact Riemannian spin manifold (M, g) , outline the evaluation of the Wodzicki residue, $\text{Wres } |\mathcal{D}|^{-n}$.

D. Regularity of a spectral triple $(\mathcal{A}, \mathcal{H}, D)$ is defined using the derivation $[|D|, \cdot]$ instead of the derivation $[D, \cdot]$. Using the standard commutative example where $\mathcal{A} = C^\infty(M)$ and $D = \mathcal{D}$ is a Dirac operator, explain why the operators $[\mathcal{D}, [\mathcal{D}, a]]$ need not be bounded for all $a \in C^\infty(M)$.

D. If $(\mathcal{A}, \mathcal{H}, D)$ is a spectral triple on an infinite-dimensional Hilbert space \mathcal{H} , explain why the operator D cannot be bounded.